

RADIATIVE CORRECTIONS TO THE DECAY $\tau \rightarrow \pi(K)\nu_\tau$

Roger Decker and Markus Finkemeier

Institut für Theoretische Teilchenphysik
Universität Karlsruhe
Kaiserstr. 12, Postfach 6980
76128 Karlsruhe
Germany

Abstract

We have calculated the $O(\alpha)$ radiative corrections to the tau decay $\tau \rightarrow \pi(K)\nu$, taking into account both the point meson contribution and the structure dependent radiation. We find for the ratio $\Gamma(\tau \rightarrow \pi\nu(\gamma))/\Gamma(\pi \rightarrow \mu\nu(\gamma))$ a radiative correction of +1.2% and for $\Gamma(\tau \rightarrow K\nu(\gamma))/\Gamma(K \rightarrow \mu\nu(\gamma))$ one of +2.0%. We compare our results with an earlier estimation and with experimental data.

Neglecting electromagnetic corrections, the decay rate $\Gamma(\tau \rightarrow M\nu_\tau)$ (where $M = \pi$ or K) can easily be predicted from $\Gamma(M \rightarrow \mu\nu_\mu)$. The relation at the order $O(\alpha^0)$ is:

$$R_{\tau/M}^{(0)} = \frac{\Gamma(\tau \rightarrow M\nu_\tau)}{\Gamma(M \rightarrow \mu\nu_\mu)} = \frac{1}{2} \frac{m_\tau^3}{m_M m_\mu^2} \frac{(1 - m_M^2/m_\tau^2)^2}{(1 - m_\mu^2/m_M^2)^2} \quad (1)$$

Using current particle data and the theoretical prediction for $\tau \rightarrow e\nu\bar{\nu}$ (for details see below), this yields the $O(\alpha^0)$ prediction

$$\left(\frac{\text{BR}(\tau \rightarrow \pi\nu) + \text{BR}(\tau \rightarrow K\nu)}{\text{BR}(\tau \rightarrow e\nu\bar{\nu})} \right)^{(theo,0)} = 0.650 \quad (2)$$

Recent experimental averages for these branching ratios have been given in Ref. [1], resulting in

$$\left(\frac{\text{BR}(\tau \rightarrow \pi\nu) + \text{BR}(\tau \rightarrow K\nu)}{\text{BR}(\tau \rightarrow e\nu\bar{\nu})} \right)^{(expt)} = 0.675 \pm 0.015 \quad (3)$$

So the $O(\alpha^0)$ theoretical prediction is off by 1.7 standard deviations. This discrepancy might either point to the importance of the neglected radiative corrections of $O(\alpha)$, or the experimental numbers might simply not be quite correct. (Note that the new τ data has greatly reduced the inconsistency between the values for the τ mass, lifetime and its leptonic branching ratio, but there is still a hint that the value given for the leptonic branching ratio might be too small.)

Marciano and Sirlin [2] have given an estimate of the QED correction. On the basis of the short distance behaviour of the weak interaction they estimate the leading $O(\alpha \ln m_Z)$ corrections to both decays as

$$R_{\tau/\pi}^{M.S.} = R_{\tau/\pi}^{(0)} \frac{1 + 2\alpha/\pi \ln(m_Z/m_\tau)}{1 + \frac{3}{2}(\alpha/\pi) \ln(m_Z/m_\pi) + \frac{1}{2}(\alpha/\pi) \ln(m_Z/m_\rho)} \quad (4)$$

(and similarly for the kaon mode, where $m_\pi \rightarrow m_K$), which, using the current particle data, yields:

$$\left(\frac{\text{BR}(\tau \rightarrow \pi\nu) + \text{BR}(\tau \rightarrow K\nu)}{\text{BR}(\tau \rightarrow e\nu\bar{\nu})} \right)^{(M.S.)} = 0.646 \quad (5)$$

This deviates from the experimental value by two standard deviations. But note that Eqn. (4) can be rewritten as

$$R_{\tau/\pi}^{M.S.} \approx R_{\tau/\pi}^{(0)} \left[1 + \frac{\alpha}{2\pi} \ln \left(\frac{m_\tau^4}{m_\pi^3 m_\rho} \right) \right] \quad (6)$$

The large mass m_Z actually drops out in the ratio and so the neglected terms which are non-leading in m_Z are of the same order of magnitude and must be taken into account in order to get a reliable result.

We conclude that the increasing experimental accuracy asks for a reconsideration of the $O(\alpha)$ corrections to these τ decays. We have calculated these corrections by taking into account both the point meson contribution and the structure dependent radiation. In the usual manner [3] we replace the vector-minus-axial vector current interaction

$$G_F \cos \theta_C f_\pi [\bar{\Psi}_l \gamma^\mu (1 - \gamma_5) \Psi_\nu] (i\partial_\mu - eA_\mu) \Phi_\pi \quad (7)$$

(which yields the point meson contribution) by the scalar-minus-pseudoscalar current interaction

$$G_F \cos \theta_C f_\pi m_l^{(0)} [\bar{\Psi}_l (1 - \gamma_5) \Psi_\nu] \Phi_\pi \quad (8)$$

where $m_l^{(0)}$ denotes the bare lepton mass.

For the structure dependent radiation we use the ansatz of Ref. [4], which we quote here for completeness (for details see there):

$$\mathcal{M}_{SD} = \frac{G_F \cos \theta_{ce}}{\sqrt{2}} \left\{ i \epsilon_{\mu\nu\rho\sigma} L^\mu \epsilon^\nu k^\rho p^\sigma \frac{F_V(t)}{m_\pi} + \bar{u}(q) \gamma_+ [(p \cdot k) \not{\epsilon} - (\epsilon \cdot p) \not{k}] u(s) \frac{F_A(t)}{m_\pi} \right\} \quad (9)$$

where the form factors are

$$\begin{aligned} F_A(t) &= -0.0116 \text{Res}_{a_1}(t) \\ F_V(t) &= \frac{-0.0270}{1.085} [\text{Res}_\rho(t) + 0.136 \text{Res}_{\rho'}(t) - 0.051 \text{Res}_{\rho''}(t)] \end{aligned} \quad (10)$$

with normalized Breit-Wigner like resonance factors

$$\text{Res}_X(t) = \frac{m_X^2}{m_X^2 - t - i m_X \Gamma_X(t)} \quad (11)$$

with energy dependent widths, as described in the quoted reference.

We write the radiative corrected decay rates as

$$\Gamma(\text{channel}) = \Gamma^0(\text{channel}) \left[1 + \frac{\alpha}{2\pi} \left(B(\text{channel}) + B_{SD}(\text{channel}) \right) \right] \quad (12)$$

where $B(\text{channel})$ is the point meson contribution, ie. the sum of virtual, soft and hard bremsstrahlung, and $B_{SD}(\text{channel})$ arises from the structure dependent radiation (and its interference with the point meson internal bremsstrahlung).

We find

$$B\left(M \rightarrow l \nu_l(\gamma)\right) = -\frac{3}{2} \Delta + \frac{3}{2} \ln \frac{m_M^2}{\mu^2} + \frac{19}{4} - \frac{2}{3} \pi^2 + 6 \ln r_l + f(r_l) \quad (13)$$

and

$$B\left(\tau \rightarrow M \nu_\tau(\gamma)\right) = -\frac{3}{2} \Delta + \frac{3}{2} \ln \frac{m_\tau^2}{\mu^2} + \frac{25}{4} - \frac{2}{3} \pi^2 + g(r_M) \quad (14)$$

where

$$\begin{aligned} f(r_l) &= 4 \left(\frac{1+r_l^2}{1-r_l^2} \ln r_l - 1 \right) \ln(1-r_l^2) - \frac{r_l^2(8-5r_l^2)}{(1-r_l^2)^2} \ln r_l \\ &\quad + 4 \frac{1+r_l^2}{1-r_l^2} \text{Li}_2(r_l^2) - \frac{r_l^2}{1-r_l^2} \left(\frac{3}{2} + \frac{4}{3} \pi^2 \right) \end{aligned} \quad (15)$$

and

$$\begin{aligned}
g(r_M) = & 4 \left(\frac{1+r_M^2}{1-r_M^2} \ln r_M - 1 \right) \ln(1-r_M^2) - \frac{r_M^2(2-5r_M^2)}{(1-r_M^2)^2} \ln r_M \\
& + 4 \frac{1+r_M^2}{1-r_M^2} \text{Li}_2(r_M^2) + \left(\frac{3}{2} - \frac{4}{3}\pi^2 \right) \frac{r_M^2}{1-r_M^2}
\end{aligned} \tag{16}$$

In these formulae Δ denotes

$$\Delta = \frac{2}{\epsilon} - \gamma_{Euler} + \ln 4\pi \tag{17}$$

where $4-\epsilon$ is the number of space-time dimensions. μ is the mass scale of dimensional regularization, $\text{Li}_2(x)$ is the dilogarithmic function

$$\text{Li}_2(x) = - \int_0^x dt \frac{\ln(1-t)}{t} \tag{18}$$

and the mass ratios r_l and r_M are defined by

$$r_l = \frac{m_l}{m_M} \qquad r_M = \frac{m_M}{m_\tau} \tag{19}$$

Note that $f(0) = g(0) = 0$, so from Eqn. (13) we find the well-known mass singularity of the total radiative correction to pion decay as $3\alpha/\pi \ln(m_l)$, whereas in the radiative correction to the tau decay the pion mass singularities cancel according to Eqn. (14). The zeroth order pion decay rate is proportional to m_l^2 , whereas the tau decay rate does not vanish for zero pion mass. So both the pion and tau results agree with the Kinoshita-Lee-Nauenburg theorem on the cancelation of mass singularities [3, 5].

For the structure dependent contribution numerical integration yields:

$$\begin{aligned}
B_{SD} \left(\pi \rightarrow \mu \nu_\mu(\gamma) \right) &= 1.8 \cdot 10^{-5} & B_{SD} \left(\pi \rightarrow e \nu_e(\gamma) \right) &= 0.66 \\
B_{SD} \left(K \rightarrow \mu \nu_\mu(\gamma) \right) &= 0.060 \\
B_{SD} \left(\tau \rightarrow \mu \nu_\mu(\gamma) \right) &= 1.46 & B_{SD} \left(\tau \rightarrow K \nu_\tau(\gamma) \right) &= 2.60
\end{aligned} \tag{20}$$

We write the radiative corrections to the ratios as

$$R_{...} = R_{...}^0 (1 + \delta R_{...}) \tag{21}$$

and find from the above

$$\begin{aligned}
\delta R_{\tau/\pi} &= +1.05\% + 0.17\% = +1.22\% \\
\delta R_{\tau/K} &= +1.67\% + 0.29\% = +1.96\%
\end{aligned} \tag{22}$$

where the first numbers (1.05 and 1.67, respectively) are the point meson contributions and the second numbers give the structure dependent part. So in both cases the model dependent contribution to the radiative correction is very small. This is true for the integrated and inclusive decay

rate, whereas in certain regions of the differential decay distribution the structure dependent part becomes dominant [4].

Note that from Eqn. (13) and (20) we can also calculate the radiative correction to the ratio of the electronic and muonic decay modes of the pion

$$R_{e/\mu} = \frac{\Gamma(\pi \rightarrow e\nu_e(\gamma))}{\Gamma(\pi \rightarrow \mu\nu_\mu(\gamma))} = \frac{m_e^2}{m_\mu^2} \left(\frac{m_\pi^2 - m_e^2}{m_\pi^2 - m_\mu^2} \right)^2 \left(1 + \delta R_{e/\mu} \right) \quad (23)$$

as

$$\delta R_{e/\mu} = -3.93\% + 0.08\% = -3.85\% \quad (24)$$

where -3.93% is the well known point pion correction [3].

From the 1992 particle data book [6] we extract

$$\begin{aligned} \Gamma(\pi \rightarrow \mu\nu_\mu(\gamma)) &= (2.528 \pm 0.002) \cdot 10^{-14} \text{ MeV} \\ \Gamma(K \rightarrow \mu\nu_\mu(\gamma)) &= (3.41 \pm 0.02) \cdot 10^{-14} \text{ MeV} \end{aligned} \quad (25)$$

and normalize to the standard model prediction for the electronic decay mode of the τ :

$$\begin{aligned} \Gamma(\tau \rightarrow e\nu\bar{\nu}(\gamma)) &= \frac{G_F^2 m_\tau^5}{192\pi^3} \left[1 - 8\frac{m_e^2}{m_\tau^2} + \frac{3}{5}\frac{m_\tau^2}{m_W^2} + \frac{\alpha}{2\pi} \left(\frac{25}{4} - \pi^2 \right) \right] \\ &= (4.033 \pm 0.005) \cdot 10^{-10} \text{ MeV} \end{aligned} \quad (26)$$

using [1]

$$m_\tau = (1777.1 \pm 0.5) \text{ MeV} \quad (27)$$

And so we arrive at our predictions:

$$\begin{aligned} \left(\frac{\text{BR}(\tau \rightarrow \pi\nu(\gamma))}{\text{BR}(\tau \rightarrow e\nu\bar{\nu}(\gamma))} \right)^{(theo, O(\alpha))} &= 0.619 \pm 0.001 \\ \left(\frac{\text{BR}(\tau \rightarrow K\nu(\gamma))}{\text{BR}(\tau \rightarrow e\nu\bar{\nu}(\gamma))} \right)^{(theo, O(\alpha))} &= 0.0410 \pm 0.0002 \\ \left(\frac{\text{BR}(\tau \rightarrow \pi\nu(\gamma)) + \text{BR}(\tau \rightarrow K\nu(\gamma))}{\text{BR}(\tau \rightarrow e\nu\bar{\nu}(\gamma))} \right)^{(theo, O(\alpha))} &= 0.660 \pm 0.001 \end{aligned} \quad (28)$$

The error bars given include only the uncertainties that result from the uncertainties of the experimental input. So our prediction agrees with the current experimental value in Eqn. (3).

We finish by comparing our approach with that of Ref. [2]. Their estimate is derived from the short distance behaviour of the weak interaction, whereas we have used a point meson and the hadronically dominated structure dependent radiation, ie. the long and medium distance behaviour. Firstly this explains the fact that the logarithms in m_Z , which our calculation gives when we put $\Delta = \ln(m_Z^2/\mu^2)$ do not agree with those in Eqn. (4). Furthermore we believe that the short distance contributions are roughly the same for tau and pion decay and so they drop out in ratios like $R_{\tau/\pi}$ at least to a good approximation. And so our calculation should give a rather reliable prediction, as it takes into account the dominant effects of these decays.

Acknowledgement

One of us (M.F.) would like to thank A. Sirlin for an illuminating discussion.

References

- [1] J.J.G. Cadenas, Experimental results on τ physics, in: Proceedings of the XIII th conference on Physics in Collision (Heidelberg, June 1993), to be published
- [2] W.J. Marciano and A. Sirlin, Phys. Rev. Lett. **61** 1815 (1988) 1815
- [3] T. Kinoshita, Phys. Rev. Lett. **2** (1959) 477
- [4] R. Decker and M. Finkemeier, “Radiative tau decays with one pseudo scalar meson”, Universität Karlsruhe Preprint, TTP93-1 (1993) (to be published in Phys. Rev. D); see also: J. Kim and L. Resnick, Phys. Rev. **D21** (1980) 1330
- [5] A. Queijeiro and J.M. Rivera, Phys. Rev. **D46** (1992) 2996
- [6] Review of Particle Properties, Phys. Rev. **D45** (1992) vol. 11